

Minneapolis Heart Institute at Abbott Northwestern Hospital Cardiovascular Laboratory Orders**TO BE COMPLETED BY PHYSICIAN**

Patient Name: _____

Diagnosis: _____

1. Select procedure to be performed:

- | | |
|--|--|
| <input type="checkbox"/> Coronary (and bypass graft if present) Angiography | <input type="checkbox"/> Oximetry Series/Shunt Study |
| <input type="checkbox"/> Left Ventriculogram | <input type="checkbox"/> Aortic Valvuloplasty |
| <input type="checkbox"/> LVEDP Only | <input type="checkbox"/> Mitral Valvuloplasty |
| <input type="checkbox"/> Possible Percutaneous Intervention (PCI) | <input type="checkbox"/> ASD/PFO Closure |
| <input type="checkbox"/> Right Heart Catheterization | <input type="checkbox"/> Ascending Aortography |
| <input type="checkbox"/> Cardiac Output | <input type="checkbox"/> Myocardial Biopsy |
| <input type="checkbox"/> Bilateral Heart Cath for Aortic Stenosis | |
| <input type="checkbox"/> Bilateral Heart Cath for Mitral Stenosis | |
| <input type="checkbox"/> Bilateral Heart Cath for Constrictive/Restrictive Disease | |
| <input type="checkbox"/> Other _____ | |

2. Other procedures/consults?
_____**3. Special scheduling instructions? (e.g. day or week, MD to perform, MD to follow in hospital, etc.)**
_____**4. Does patient have a history of allergy to iodinated contrast agents? YES NO**

Pre-medication regimen: (see preprinted prescription)
Prednisone 60 mg night before procedure
Prednisone 60 mg morning of procedure
(Antihistamines can be given after arrival in Prep/Recovery)

5. Is the patient diabetic? YES NO (instructions in the packet)**6. Will the patient need pre-medication with Mucomyst? YES NO**

Acetylcysteine/Mucomyst regimen: (see preprinted prescription) 600 mg for 2 doses the day before procedure and 2 doses the day of procedure

7. Does the patient take Coumadin®? YES NO

If yes, instruct the patient to stop 3 (or more if desired) days prior to procedure

8. Does the patient take Pradaxa? YES NO

If yes, what is the patient's GFR?

- If GFR > 50 direct the patient not to take any Pradaxa the day before or day of the procedure.
- If GFR ≤ 50 instruct the patient not to take any Pradaxa two days before the day of the procedure.

Signature: _____